



[Home](#) ▶ [All issues](#) ▶ [Volume 403 / No 1 \(May III 2003\)](#) ▶ [A&A, 403 1 \(2003\) L19-L23](#) ▶ [Abstract](#)

Free Access

Issue	A&A Volume 403, Number 1, May III 2003
Page(s)	L19 - L23
Section	Letters
DOI	https://doi.org/10.1051/0004-6361:20030450
Published online	29 April 2003

A&A 403, L19-L23 (2003)

Letter to the Editor

A broad absorption feature in the X-ray spectrum of the isolated neutron star RBS1223 (1RXS J130848.6+212708)*

F. Haberl¹, A. D. Schwope², V. Hambaryan², G. Hasinger¹ and C. Motch³

¹ Max-Planck-Institut für extraterrestrische Physik, Giessenbachstraße, 85748 Garching, Germany

² Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

³ Observatoire Astronomique, UA 1280 CNRS, 11 rue de l'Université, 67000 Strasbourg, France

Corresponding author: F. Haberl, fwh@mpe.mpg.de

Received: 6 March 2003

Accepted: 25 March 2003

Abstract

X-ray spectra of the isolated neutron star RBS1223 obtained with the instruments on board XMM-Newton in December 2001 and January 2003 show deviations from a Planckian energy distribution at energies below 500 eV. The spectra are well fit when a broad, Gaussian-shaped absorption line with $\sigma = 100$ eV and centered at an energy of 300 eV is added to an absorbed blackbody model. The resulting equivalent width of the line is ~ 150 eV. However, the spectral resolution at these low energies of the EPIC detectors and the lower statistical quality and restricted energy band of the RGS instruments are not sufficient to exclude even broader lines at energies down to 100 eV or several unresolved lines. The most likely interpretation of the absorption feature is a cyclotron absorption line produced by protons in the magnetic field of the neutron star. In this picture line energies of 100–300 eV yield a magnetic field strength of $2\text{--}6 \times 10^{13}$ G for a neutron star with canonical mass and radius. Folding light curves from different energy bands at a period of 10.31 s, which implies a double peaked pulse profile, shows different hardness ratios for the two peaks. This confirms that the true spin period of is twice as long as originally thought and suggests variations in cyclotron absorption with pulse phase. We also propose that changes in photo-electric absorption seen in phase resolved spectra of by Cropper et al. ([CITE]), when formally fit with an absorbed blackbody model, are caused instead by cyclotron absorption varying with pulse phase.

* Based on observations with XMM-Newton, an ESA Science Mission with instruments and contributions directly funded by ESA Member states and the USA (NASA).

© ESO, 2003

Table of Contents

Article

	Full	PDF (219
Abstract	HTML	KB)
References	Simbad Objects	
NASA ADS Abstract		
Service		

Metrics

Show article metrics

Services

Articles citing this article

CrossRef (87)

Same authors

- Google Scholar
- NASA ADS
- EDP Sciences database

Recommend this article

Send to my Kindle
Download citation

Related Articles

A phase-dependent absorption line in the spectrum of the X-ray pulsar RX J0720.4-3125

A&A 419, 1077-1085 (2004)

Evidence for precession of the isolated neutron star RX J0720.4-3125

A&A 451, L17-L21 (2006)

XMM-Newton RGS spectrum of RX J0720.4-3125: an absorption feature at 0.57 keV

A&A 497, L9-L12 (2009)

[More](#)

Bookmarking



Astronomy & Astrophysics (A&A)

Editor-in-Chief: T. Forveille

Letters Editor-in-Chief: J. Alves

Managing Editor: D. Elbaz

ISSN: 0004-6361 ; e-ISSN: 1432-0746

Frequency: 12 volumes per year

Published by: EDP Sciences

Mirror sites: [CDS](#) | [EDP Sciences](#)

© The European Southern Observatory ([ESO](#))

