

# Prey of killer whales (*Orcinus orca*) in Iceland

PLOS ONE

PUBLISH

ABOUT

BROWSE

SEARCH



advanced search

OPEN ACCESS PEER-REVIEWED

RESEARCH ARTICLE

0 Save

5 Citation

## Prey of killer whales (*Orcinus orca*) in Iceland

Filipa I. P. Samarra  Manuela Bassoi, Julie Béseau, Margrét Ó. Elíasdóttir, Karl Gunnarsson, Marie-Thérèse Mrusczo, Marianne Rasmussen, Jonathan N. Rempel, Baldur Thorvaldsson, Gísli A. Víkingsson

4,567 View

3 Share

Published: December 12, 2018 • <https://doi.org/10.1371/journal.pone.0207287>

Article	Authors	Metrics	Comments	Media Coverage
---------	---------	---------	----------	----------------


Download PDF

Print

Share

- Reader Comments (0)
- Media Coverage
- Figures

### Figures



ADVERTISEMENT

### Abstract

Killer whales have a cosmopolitan distribution and as a species are generalists, feeding on a variety of prey. However, local populations tend to specialise on specific prey types. In Icelandic waters, killer whales are generally associated with herring and, thus, have been presumed to be herring specialists. However, recent studies suggest a more complex foraging ecology, possibly including a mosaic of strategies. With increased observational effort in recent years due to research and whale-watching activities, there have been several reports of interactions with different prey, including confirmed predation events. In this study we aimed to summarise the range of potential prey of killer whales observed in Icelandic waters. We report on 12 previously unpublished accounts and review 15 accounts published in the scientific literature or local newspapers, making a total of 27 events where killer whales were observed interacting with actual or potential prey. Thirteen different species, including birds ( $n = 1$ ), cephalopods ( $n = 1$ ), fish ( $n = 5$ ) and marine mammals ( $n = 6$ ), are reported, although herring is by far the species that killer whales are most often observed interacting with. This study provides the first summary of actual and suspected killer whale prey in Icelandic waters, and contributes towards our understanding of this population's prey preferences. However, describing the diet of individuals/groups was not possible and this study points to a need for continued monitoring to understand the intricacies of killer whale foraging behaviour in this area.

**Citation:** Samarra FIP, Bassoi M, Béseau J, Elíasdóttir MÓ, Gunnarsson K, Mrusczo M-T, et al. (2018) Prey of killer whales (*Orcinus orca*) in Iceland. PLoS ONE 13(12): e0207287. <https://doi.org/10.1371/journal.pone.0207287>

**Editor:** David Hyrenbach, Hawaii Pacific University, UNITED STATES

**Received:** June 21, 2018; **Accepted:** October 29, 2018; **Published:** December 12, 2018

**Copyright:** © 2018 Samarra et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** All relevant data are within the paper.

**Funding:** MB was employed by Elding Whale Watching, MOE was employed by Whales Hauganes ehf., MTM was employed by Láki Tours, JNR was employed by Special Tours and BT was employed by Iceland Pro Cruises. The funders provided support in the form of salaries for authors and in all cases (except Iceland Pro Cruises) provided platforms from where observations were made, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section. Funding for dedicated research was provided by the Icelandic Research Fund. The funder had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** MB was employed by Elding Whale Watching, MOE was employed by Whales Hauganes ehf., MTM was employed by Láki Tours, JNR was employed by Special Tours and BT was employed by Iceland Pro Cruises. These commercial affiliations do not alter our adherence to PLOS ONE policies on sharing data and materials.

## Introduction

Killer whales (*Orcinus orca*, Linnaeus, 1758), as a species, have been reported to predate on numerous prey including fish, marine mammals, cephalopods, marine turtles and birds [1, 2]. However, several populations specialize on particular prey types and ecotypes diverging in diet, morphology, genetics and behaviour have been recognised in the North Pacific and Antarctica (e.g. [3–9]). Nevertheless, killer whales with an apparent mixed diet that includes fish and marine mammals have been reported in several locations [10, 11]. Foraging ecology is tightly related to the social organization, life history and behaviour of killer whales [12] and, ultimately, may play a major role in the social segregation and genetic divergence of populations [13, 14]. Understanding the foraging ecology of each population is relevant to assess both their conservation status and role in marine ecosystems.

In the Northeast Atlantic, two killer whale ecotypes have been suggested based on analyses of morphological traits, stable isotope ratios and tooth wear from museum and stranded specimen; a generalist Type 1, that presumably feeds upon fish and to some extent seals, and a marine mammal specialist Type 2 [15]. High variance in nitrogen stable isotope ratios within Type 1 suggested individual and/or group variation in the proportion of prey types consumed [15]. A subsequent genetic study identified three significantly differentiated populations overlapping spatially with the distribution of Atlantic herring (*Clupea harengus*, Linnaeus, 1758), Atlantic mackerel (*Scomber scombrus*, Linnaeus, 1758) and Atlantic bluefin tuna (*Thunnus thynnus*, Linnaeus, 1758), on which some members of each population had been seen predated [16]. Population A included herring-feeding individuals that were sampled within the ranges of the Norwegian, Icelandic and North Sea herring stocks. Population B included individuals sampled from the North Sea to the West coast of Iceland that appeared to spatially associate with the distribution of the Northeast Atlantic mackerel stock, including one individual sampled from a trawler fishing boat whilst feeding on mackerel discards. Finally, population C was composed solely of individuals sampled in the Canary Islands and in the Strait of Gibraltar, the latter known to target tuna (e.g., [17]).

Killer whales have long been known to occur around Iceland, particularly on seasonal herring grounds, where killer whales are observed feeding upon herring [18]. Although initially suggested to be a population of herring-specialists [18], genetic studies suggested that killer whales sampled in Icelandic waters belong to two separate populations: population A, which associates with herring and population B, which is thought to associate with mackerel [16]. Furthermore, recent studies showed that killer whales occurring in Icelandic coastal waters have different movement and foraging strategies. While some individuals appear to specialise on herring year-round, others prey on it only seasonally or even opportunistically and include other unknown, higher trophic level prey in their diet [19]. Indeed, a few individuals occur regularly in herring overwintering grounds in Iceland but in spring and summer travel to Scotland where they have been seen feeding upon seals [19–21]. Early reports of feeding upon marine mammals and seabirds [22] indicated that prey targeted by killer whales in Iceland extended beyond herring. Nevertheless, target prey species remained relatively unknown. Thus, there is still little understanding of the prey preferences of Icelandic killer whales as well as the potential existence of different ecotypes in these waters.


In recent years, there has been an increase in whale-watching in Iceland, which now takes place

in several locations across the country [23]. This has led to increased numbers of killer whale encounters outside herring grounds and interactions with other potential prey have been reported more frequently. In this study, we report on recent unpublished observations of killer whale interactions with potential prey and review available information from published sources to provide a summary of potential prey targeted by killer whales in Icelandic waters. This study aimed to increase knowledge of the prey of Icelandic killer whales, to contribute towards an assessment of the species' role within this ecosystem.

## Materials and methods

All observations from whale-watching vessels were conducted in accordance with local whale watching guidelines. All field research was carried out in compliance with local regulations and under an institutional permit for the Marine and Freshwater Research Institute, Reykjavík.

Sightings of killer whales interacting with various species in Icelandic waters were compiled from several sources, including previously unpublished reports from dedicated research and commercial whale-watching operations, as well as published accounts on newspapers or other scientific and non-scientific published literature. All observations were summarised in [Table 1](#). Previously published reports are briefly mentioned in the results, along with detailed accounts of recent, unpublished observations. The reported times of events are all in local time (UTC). Observations were conducted opportunistically onboard whale-watching boats or from dedicated research vessels. When available, photographs were used to identify the individual killer whales involved in each incident. Individuals were compared with an existing photo-identification catalogue [24], to understand if those individuals had been observed consuming other prey before.



**Table 1. Summary of interactions between killer whale and other species in Icelandic waters.**


Prey is divided into four types: birds, cephalopods, fish and marine mammals. Date is given as dd/mm/yyyy when available or by the month and year. When available, information on location of events is provided as approximate locations or exact coordinates. Approximate and exact locations are plotted in [Fig 1](#), unless the location information provided was too inaccurate, such as "East of Iceland" (see events #15 and 16).

<https://doi.org/10.1371/journal.pone.0207287.t001>

Published records of interactions between killer whales and other species were obtained from local printed media accessed through a public database (timarit.is). The keyword 'háhyrningur' (Icelandic common name for killer whale) and all its possible inflections were used to search the database. The resulting news items included material published between 1880 and 2016 and were investigated to find relevant accounts of killer whales confirmed interacting with other species. Relevant interactions are listed in [Table 1](#) with references to the newspaper and date of publication. Accounts published in the scientific literature were also summarised. Only interactions where killer whales were observed chasing, harassing and attacking or presumably attacking other species, defined as predatory interactions [2], were included. Additional observations of stomach contents or prey handling where the capture was not recorded were also included.

## Results

A total of 13 different potential prey species were identified in this study, including birds, cephalopods, fish and marine mammals ([Table 1](#)). Predation events occurred all around Iceland ([Fig 1](#)). Twelve recently observed accounts of killer whale interactions with different prey are reported here for the first time ([Table 1](#)). The remaining 15 accounts reviewed consisted of 3 events reported previously in the scientific literature and 12 events reported in newspaper articles. Predation and lethal harassment can be difficult to distinguish, particularly from popular accounts, so we included a description in [Table 1](#) of what was in the published literature, without assuming consumption unless clearly observed. For all the previously unpublished observations reported here, we state when consumption of prey was not confirmed. A detailed description of previously unpublished accounts, along with a mention of accounts previously reported in the literature, is given below, separated into four main species groups: 1) birds; 2) cephalopods; 3) fish; and 4) marine mammals.



Download:

PPT PowerPoint slide

PNG larger image

TIFF original image

**Fig 1. Map of approximate locations of events of killer whale interactions with the different prey species described in this study.**

Numbers correspond to events listed in [Table 1](#). If more than one event occurred in the same location the numbers are listed vertically.

<https://doi.org/10.1371/journal.pone.0207287.g001>

## Birds

### Common eider (*Somateria mollissima*, Linnaeus, 1758).

In February 1986, a group of approximately 12 to 15 killer whales, including two adult males, were observed in the inner part of Berufjörður (E Iceland), in Fossárvík ([Table 1](#), event #1). Two relatively small animals were observed surfacing where a flock of common eiders was densely grouped close to the shore. One of the whales grabbed a male eider and took it underwater and then the rest of the eider flock fled into a cove. The bird was not seen again so it was presumed to be consumed. The killer whales disappeared underwater and then came up under the birds only a few meters away from the observers. The eiders all rushed away just before the whales surfaced. Shortly after, the killer whales left the area to join the rest of the group out on the fjord. No photo-identification data were available for this event.

On 18 April 1998, a small group of killer whales was observed hunting, attacking and eating common eiders along the coast of Reykjavík ([Table 1](#), #2). The killer whale group was composed of two adult males, a female with a calf and one subadult male/adult female. The killer whales moved systematically along the coast (approximately 10–100 m from the shore) and surfaced repeatedly in groups of eiders. At least two eiders were observed swallowed and probably more were eaten. Photographs of only the right sides of the whales involved in this event were available but it was not possible to identify any individuals.

Three further accounts of predation on common eiders were reported in newspapers and occurred in Ísafjarðardjúp (NW Iceland) in 2003 ([Table 1](#), #3), Djúpvogur (E Iceland) in 2006 ([Table 1](#), #4) and Berufjörður (E Iceland) in 2010 ([Table 1](#), #5).

On 16 November 2014 at approximately 14:33, two killer whales (identified as IS015 and IS229, [24]) were observed attacking common eiders just outside Grundarfjörður ([Table 1](#), #6). Both individuals were observed lunging out of the water in the locations where eiders were sitting at the sea surface. On two occasions, the bird targeted seemed to have been pulled under the surface. Prey consumption was not confirmed.

## Cephalopods

Predation on cephalopods was reported once in the scientific literature, as part of a list of items found in the stomach contents of killer whales caught by whalers ([Table 1](#), #7).

## Fish

### Herring (*Clupea harengus*, Linnaeus, 1758).

There are numerous observations of killer whales occurring in seasonal herring grounds, making this by far the prey that killer whales are most commonly observed consuming in Iceland. The first published scientific book mentioning herring as a main prey of killer whales dates back to 1932 and since then there have been numerous studies reporting herring consumption ([Table 1](#), #8). Indeed, killer whales in Iceland were proposed to be herring-specialists [18], which has been supported by recent studies [19, 21].

### Lumpfish (*Cyclopterus lumpus*, Linnaeus, 1758).

Three killer whales (identified as IS015, IS113 and IS229, [24]) were observed in the morning of the 3 March 2014, at approximately 11:13, just outside Kolgrafafjörður ([Table 1](#), #9). The whales were tracked from a whale-watching boat following the coastline, heading east. Approximately 1h later, at the entrance of the fjord, individual IS229 was observed with a fish in its mouth ([Fig 2](#)). Photographs were later used for identification of the fish species, consulting experts that confirmed it to be a lumpfish. Great Black-backed gulls (*Larus marinus*) and Glaucous gulls (*Larus hyperboreus*) followed the killer whale and attempted to scavenge on pieces of fish from the surface, but consumption of the fish by the whale was not confirmed. A few minutes later, the whale-watching boat departed the area. A dedicated research vessel found the same individuals in that area in the afternoon. At approximately 13:44, one whale was again seen with the head out of the water and holding something in its mouth and once again it was followed by gulls that picked up a piece of what appeared to be fish. However, it was not possible to determine the

species on this occasion. The killer whales were then observed travelling very close to a seal haul-out location, closely following the shoreline. However, no interactions were observed even though at least one seal was present in the water.



On 11 June 2016, one animal was observed at approximately 15:36 just off the tip of the Snæfellsnes peninsula (Table 1, #10). The whale was observed lunging often and spyhopped once, followed by a backflip while tossing a lumpfish up with its rostrum. The individual was then seen rolling on its side, no further surface activity was observed and the individual was just milling. Prey consumption was not confirmed. Another five individuals were present in the area but were not interacting with fish. Photographs from this event were available but it was not possible to match the individual involved.

### Salmon (*Salmo salar*, Linnaeus, 1758).

On 15 July 2015, at approximately 16:12, a dedicated research vessel was following a group of four killer whales, including two adult females, a calf and one juvenile in Vestmannaeyjar (Table 1, #11). The group was observed swimming erratically at the surface turning sharply as if chasing something underwater. This went on for a few minutes until the whales were seen tossing a fish in the air (Fig 3). Photographs of the event were being collected throughout for photo-identification. Although the fish was observed being tossed a few times, consumption could not be confirmed due to the distance to the whales. Later inspection of the photographs with consultation of experts revealed the fish species to be salmon (*Salmo salar*).



Atlantic halibut (*Hippoglossus hippoglossus*, Linnaeus, 1758) and Greenland halibut (*Reinhardtius hippoglossoides*, Walbaum, 1972).

Accounts from newspapers reported depredation from longline fisheries for Atlantic halibut in 1978 (Table 1, #12), 1979 (Table 1, #13) and 1992 (Table 1, #14), and depredation from longline fisheries for Greenland halibut in 1970 (Table 1, #15) and 1971 (Table 1, #16).

### Marine mammals

Pilot whales (*Globicephala melas*, Traill, 1809).

Possible predation on long-finned pilot whales was reported once in the scientific literature (Table 1, #17).

Common minke whale (*Balaenoptera acutorostrata*, Lacepede, 1804).

Accounts of predation on minke whales, including confirmed consumption, were published in newspapers in 1915 (Table 1, #18) and 1997 (Table 1, #19).

On 22 July 2008 killer whales were observed in Skjálfandi bay off Húsavík (N Iceland; Table 1, #20). The sighting started with the observation of a common minke whale jumping out of the water. A whale-watching vessel headed in that direction to find a group of killer whales chasing the minke whale (Fig 4). The group consisted of approximately 20 individuals of all age/sex classes including one adult male. Two to four killer whales took turns surrounding the minke whale, but the adult male always stayed at a distance of more than 50 m and did not take part in

the attack itself. The minke whale was observed jumping out of the water several times. Towards the end of the attack, the minke whale's tongue was hanging out of its mouth and the killer whales dragged it underwater and kept it under the surface. The encounter lasted approximately 45 minutes, but consumption was not confirmed. The killer whales left the bay once the minke whale was dead. Photographs of some of the whales involved in this event were available but it was not possible to identify the individuals involved.



Harbour porpoise (*Phocoena phocoena*, Linnaeus, 1758).

Accounts of predation on harbour porpoises were published in newspapers in 1982 ( [Table 1, #21](#)) and 1997 ([Table 1, #19](#)).

On 27 June 2012 a group of about five killer whales, including one adult male, was observed by whale-watching vessels attacking a harbour porpoise as well as a harbour seal (*Phoca vitulina*) in Skjálfandi Bay off Húsavík ([Table 1, #22](#)) within one hour. One killer whale was observed taking the harbour porpoise in its mouth, but consumption of both prey was not confirmed. The whales left the area soon after the hunt ended. Photographs from this event were available but not suitable to confirm matching for photo-identification.

On 23 September 2017, at approximately 10:05 a group of killer whales was seen travelling in Eyjafjörður ([Table 1, #23](#)). The group was estimated as 12 individuals including two adult males and a young juvenile. Two humpback whales (*Megaptera novaeangliae*, Borowski, 1781) were also seen approximately 700 m away, nearer the coast, and a group of approximately 10–15 harbour porpoises was in the same general area, nearer the middle of the fjord. The killer whales travelled initially in a southern direction into the fjord, but then turned sharply and headed back to the area where the harbour porpoises were. Four to six killer whales, including an adult male, started speeding towards the harbour porpoises where they ended up making a kill ([Fig 5](#)). Following the kill, all whales seemed to join and consume the prey. At least one porpoise was killed, but the actual number taken is unknown. The killer whales then left the fjord, remaining close to the coastline. The encounter lasted approximately 50 minutes. Photographs of some of the whales involved in this event were available but it was not possible to identify the individuals involved.



White-beaked dolphin (*Lagenorhynchus albirostris*, Gray, 1846).

On 17 December 2017, at approximately 15:30, a single white-beaked dolphin was observed from a whale-watching boat in Faxaflói Bay ([Table 1, #24](#)). This was followed by a sighting of a killer whale calf leaping out of the water nearby. Soon the rest of the killer whale group, making a total of five whales, appeared and surrounded the dolphin, with a lot of water splashing at the surface. The dolphin surfaced a few more times, then disappeared. Following the sighting of one of the adult whales with a large white 'chunk' in its mouth, the dolphin was presumed to have been successfully hunted ([Fig 6](#)). The whales then milled around the boat for some time. The encounter lasted approximately 25 minutes, and then the boat left the group. Photographs from this event were available but not of suitable quality to enable photo-identification.



Download:

PPT PowerPoint slide

PNG larger image

TIFF original image

**Fig 6. Killer whale predation on a white-beaked dolphin.**

Observed in Faxaflói bay (SW Iceland) on 17 December 2017 (photograph by J. Rempel):

A) killer whale ramming the ventral side of the dolphin; B) killer whale holding a white piece of presumed dolphin blubber in its mouth.

<https://doi.org/10.1371/journal.pone.0207287.g006>

Grey seal (*Halichoerus grypus*, Fabricius, 1791).

Predation on grey seals was published in newspapers in 1967 ( [Table 1](#), #25).

On 7 July 1993, a group of seven killer whales including at least one adult male and one presumed calf, was sighted circling the shallows of Flóta-flögur in Breiðafjörður by observers on a small fishing boat ([Fig 7](#); [Table 1](#), #26). As a grey seal surfaced in the middle of the group, the killer whales increased their swimming speed and swam in a tighter circle. The seal then dove into an underwater kelp forest while the whales kept circling the shoals. After several surfacings by the seal, the whales took turns in attacking the seal, repeatedly ramming it and hitting it with their fluke. The whales repeatedly rammed and hit the seal and the seal could be seen clearly bleeding. The captain then took his shotgun and shot the seal when it surfaced. When the captain moved the boat towards the seal to try to hook it out of the water, one of the whales came alongside the boat from behind and grabbed the seal and swam off with it. The group joined in and a large red patch appeared on the sea surface. The event lasted for an estimated one hour and photographs were available but not of suitable quality for photo-identification.



Download:

PPT PowerPoint slide

PNG larger image

TIFF original image

**Fig 7. Killer whale predation on a grey seal.**

Observed in Breiðafjörður (W Iceland) on 7 July 1993 (photograph by K. Gunnarsson).

<https://doi.org/10.1371/journal.pone.0207287.g007>

Harbour seal (*Phoca vitulina*, Linnaeus, 1758).

Predation on a harbour seal, in the same event of predation on a harbour porpoise, was observed in Skjálfandi bay in 2012 (see above; [Table 1](#), #27).

## Discussion

In this study we present records of 13 different species thought to be prey to killer whales occurring in Icelandic waters, including birds, cephalopods, fish and marine mammals. This list includes many but not all of the same prey reported in neighbouring regions of the North Atlantic ([Table 2](#)). All species except lumpfish, Atlantic halibut and Greenland halibut have been reported as killer whale prey in neighbouring regions. In contrast, a total of 13 species including birds (n = 5), jellyfish, fish (n = 3) and marine mammals (n = 4) have been reported in neighbouring regions but not in Iceland to date. The strandings network managed by the Marine and Freshwater Research Institute includes very few stomach contents from stranded whales and these have not been analysed yet. In addition, sub-surface consumption of fish, cephalopods or other prey is often difficult to determine. Thus, it is not unlikely that future work will increase the list of known prey in Icelandic waters.



Download:

PPT PowerPoint slide

PNG larger image

TIFF original image

**Table 2. Summary of potential and actual prey items reported for killer whales in Iceland (this study) and a comparison with some neighbouring regions.**

The only bird species confirmed as prey of killer whales was the common eider. Eiders are very common in Iceland, with an estimated population size of ~850,000 and have a wide distribution [58], thus could be a reliable prey resource for killer whales. Common eider predation appears to occur regularly and is widespread, thus killer whales may be a significant predator of this species in Icelandic waters. However, it is unlikely killer whales rely heavily on this prey as a main source of food. Indeed, killer whales preying on birds may do so for “play” [59] or to help develop skills for hunting and handling prey [3].

Cephalopods are presumed to be an underappreciated component of the diet of transient killer whales and closely related ecotypes in the North Pacific Ocean [60]. We could only find one record of squid consumption based on stomach contents of a killer whale caught in the offshore waters east of Iceland [28]. Many records of cephalopod consumption by killer whales come from caught or stranded specimens where stomach contents have been collected [28, 60], indicating that visual observations of this behaviour are scarce, probably because they would take place at depth. Thus, it is possible that this behaviour occurs but simply has not been observed in Iceland.

Lumpfish, salmon, Atlantic halibut and Greenland halibut were the only fish species, other than the well-known herring prey, to be observed, although consumption was not confirmed in some cases. One of the interactions with lumpfish was by the same whales involved in an interaction with a common eider, suggesting these whales may target diverse prey. These killer whales are known to occur in Iceland in the winter apparently feeding upon herring but then travel to Scotland where they are regularly sighted in the spring and summer [20]. This indicates they do not specialize on the Icelandic summer-spawning herring stock year-round. Indeed, stable isotope measurements of IS015 and IS229 revealed inclusion of prey other than herring in their diet, suggesting these individuals have a varied diet relative to putative herring-specialists [19]. Lumpfish migrate from offshore feeding areas to coastal areas during the breeding season (approximately March to August) [61] and, thus, could be a prey seasonally available. In Iceland, lumpfish is also known to be an important part of the diet of sperm whales (*Physeter macrocephalus*, Linnaeus, 1758) [62].

One predation event involved the chasing and capture of a salmon in Vestmannaeyjar by individuals IS130, IS340, IS412 and a calf. These whales are regularly seen in the area and are believed to follow herring year-round because they are sighted in both herring overwintering and spawning grounds [21]. Vestmannaeyjar is a herring summer spawning ground [63], but also lies in the path of the Atlantic salmon’s yearly migration to the Rangá, Þjórsá and Ölfusá rivers in the south of Iceland. From spring to late summer, tens of thousands of salmon appear in these rivers, with estimates for the year of 2013 of around 12,000 salmon migrating to the Þjórsá river alone [64]. Although over several field seasons in Vestmannaeyjar this was the only confirmed event involving salmon, it cannot be excluded that this may be a prey targeted regularly. For example, for the majority of observed feeding events of ‘resident’ killer whales off the Northwest Pacific prey identification was achieved through sampling of tissues and scales rather than by direct observation of prey species [65]. At least 96% of these events were of predation on salmonids.

Atlantic halibut and Greenland halibut depredation from longlines were reported in newspapers, including accounts that fishermen had to give up the longline fisheries due to depredation by killer whales. Indeed, in 2002 a fisherman reported that the Greenland halibut fishery changed fishing gear to nets because using longlines failed due to depredation by killer whales and sperm whales in the east and west of Iceland, respectively [66]. The Atlantic halibut population in Icelandic waters decreased drastically during 1985–1992 and has been protected from direct fishing since 2012 [67]. On the other hand, Greenland halibut is one of the most abundant groundfish species inhabiting the waters west, north and east of Iceland [68]. Because the Greenland halibut fishery is nowadays largely undertaken by trawlers, with only some vessels using gillnet and longline [69], depredation may no longer be an issue at present.

As for marine mammals, a total of six prey species were reported in this study. Long-finned pilot whales have been previously recorded in both predatory and non-predatory interactions with killer whales [2]. In Iceland, we could find only one record of killer whales attacking pilot whales. In fact, interactions between pilot whales and killer whales have occurred often in recent years, but those are of pilot whales chasing killer whales off Vestmannaeyjar (Samarra, personal observation), as observed in Gibraltar [70] and Norway [71]. This suggests killer whale predation on pilot whales is not common in Iceland, however the interspecific dynamics of these species warrant further study.

In a study reviewing interactions of killer whales with marine mammal prey, minke whales were recorded as one of the most common cetacean prey species [2]. In addition to accounts of minke whale predation in local newspapers, in this study we report one incidence of an attack on a minke whale in Skjálfandi Bay where consumption was not confirmed. Surplus killing by killer whales, whereby prey is killed but not consumed, has been previously observed in various regions (see [2]) but the reasons for this behaviour are unknown [2]. The observation in Skjálfandi Bay described in this study resembles those of coordinated attacks by transient killer whales on minke whales in the Pacific, involving prolonged chases and ending in ramming and/or asphyxiation [72]. Further reports [22] and tooth rake marks observed on the skin of minke



whales [73] suggest attacks on minke whales are likely to be common in Icelandic waters.

Harbour porpoises are commonly taken by killer whales in several locations [2]. The reports in this study confirm that predation on harbour porpoises also occurs in Iceland. The harbour porpoise is one of the most common small cetaceans in coastal Icelandic waters, however there is little information on population size and conservation status [74]. In this study we included four reports of harbour porpoise predation. However, the extent to which killer whales may predate on this species remains unknown. The fact that the killer whales involved in one of the predation events reported here were not matched to an existing photo-identification catalogue could suggest the occurrence of infrequently seen individuals that may belong to a separate, marine-mammal specialist ecotype. However, further predation records accompanied with photo-identification data will be necessary to assess this possibility.

Killer whales and white-beaked dolphins have been seen in non-predatory interactions in both Scotland and Iceland [2], and in Iceland the two species have been seen feeding together [18]. Killer whales have been observed hunting or attacking white-beaked dolphins in Scotland [53, 54] but detailed accounts of these events were not provided. In Iceland, skin lesions consistent with killer whale rake marks in white-beaked dolphins from Skjálfandi Bay and Faxaflói Bay suggested that killer whales are natural predators [73]. However, the rake mark prevalence was low [73, 75], indicating that interactions with killer whales may only occur occasionally. Our study reports the first observation of confirmed predation on white-beaked dolphins by killer whales in Iceland, but based on a single event supports a low prevalence of predation on this species.

Several seal species have been targeted by killer whales worldwide. Harbour seals appear to be one of the main seal prey [2], with some killer whales apparently specialising on this prey [76]. Only grey and harbour seals occur regularly throughout the year and breed in Iceland [77]. Both seal populations have wide distributions but have been declining in recent decades [77, 78]. The factors leading to population declines in Iceland are poorly understood, although hunting and bycatch, as well as prey availability are thought to be main contributors [78]. In this study, we included confirmed predation events on both seal species. Killer whales are thought to be the main natural predator of seals in Iceland [77] and further anecdotal evidence suggests that predation on seals occurs more often than suggested by the few accounts presented here. However, assessing the predation pressure by killer whales on both seal species warrants further study.

This study shows that killer whales in Iceland take a wide range of prey. However, the extent of predation pressure on these prey species remains unknown and other potential prey items could remain undetected. Rake marks suggest interactions with other potential prey. For example, humpback whales sighted in Iceland exhibited a scarring rate of approximately 8% [79]. Predatory interactions are thought to occur in high-latitude waters [79], suggesting some of these events could happen in Iceland. A separate study reported a rake mark frequency in humpback whales of 13% in Icelandic waters and showed that generally predation of humpback whales by killer whales was largely confined to young animals [80], as seen in other locations [81]. However, to our knowledge, predatory interactions between humpback whales and killer whales in Iceland have not been documented to date. Tooth rake and bite marks have also frequently been observed on fin (*Balaenoptera physalus*, Linnaeus, 1758) and sei (*Balaenoptera borealis*, Lesson, 1828) whales in Icelandic waters (G. Víkingsson, personal observation). Although some killer whales that had stranded in Iceland grouped genetically with whales spatially associating and feeding upon mackerel [16], to date we have not found observations of mackerel predation. Thus, with an increased observer effort in the coastal waters of Iceland, the list of potential prey species may increase in the future.

The killer whale population in Iceland shows diversity in foraging strategies, with some whales apparently specialising upon herring while others include higher trophic level prey in their diet [19]. This is consistent with the existence of an ecological gradient where individuals and/or groups vary in the proportion of different prey items consumed, as suggested for Type 1 killer whales [15]. Given our observations of interactions with marine mammals, we cannot exclude the possibility that marine mammal specialist killer whales [15] occur in Icelandic waters. Some of the whales observed off Húsavík attacking a minke whale had narrow and fainter coloured saddle patches, which were similar to those described in other regions (Hawaii [82]; Caribbean [83]; Gulf of Mexico [84]) but unlike most saddle patches in an existing photo-identification catalogue [24]. However, this feature was not shared among all individuals involved in the event suggesting they were not of a single morphotype. Killer whales with diets presumed to include herring and other high trophic level prey were observed interacting with a lumpfish and a common eider, supporting a diverse diet in these individuals. On the other hand, killer whales believed to feed predominantly on herring were observed interacting with salmon. Prey specialisation does not imply exclusivity, as seen by 'resident' killer whales in the NE Pacific that specialise on salmon but occasionally take other fish [3]. However, it is also known that killer whales often kill prey they do not intend to eat [59] and we cannot rule out the possibility that it occurred in some of the events we describe. Clearly, continued observations will be essential to assess the frequency by which different prey items are consumed and evaluate their contribution to the diet of individuals/groups observed in Iceland.

## Acknowledgments

We would like to sincerely thank the following whale-watching tour operators for providing platforms from where sightings were collected: Special Tours in Faxaflói bay, Láki Tours in

Breidafjörður, Elding and Whale Watching Hauganes in Eyjafjörður and North Sailing and Gentle Giants in Húsavík. We would like to sincerely thank Benóný Jónsson and Einar Jónsson for providing information about salmon and cephalopod occurrence in Iceland, respectively. Finally, we would like to thank Wouter Jan Strietman for providing the photograph of lumpfish predation by killer whales, Paul Wensveen for helpful comments on the manuscript, as well as Dr. R. Pitman, E. Jourdain and the editor for very useful comments that improved the manuscript.

## References

1. Hoyt E. Orca, the whale called killer. Ontario: Camden House Publishing, 1984.
2. Jefferson TA, Stacey PJ, Baird RW. A review of killer whale interactions with other marine mammals: predation to co-existence. *Mamm Rev.* 1991;21: 151–180.  
[View Article](#) • [Google Scholar](#)
3. Ford JKB, Ellis GM, Barrett-Lennard LG, Morton AB, Palm RS, Balcomb KC III. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Can J Zool.* 1998;76: 1456–1471.  
[View Article](#) • [Google Scholar](#)
4. Pitman RL, Ensor P. Three forms of killer whales (*Orcinus orca*) in Antarctic waters. *J Cetacean Res Manag.* 2003;5: 131–139.  
[View Article](#) • [Google Scholar](#)
5. Pitman RL, Perryman WL, LeRoi D, Eilers E. A dwarf form of killer whale in Antarctica. *J Mamm.* 2007;88: 43–48.  
[View Article](#) • [Google Scholar](#)
6. Pitman RL, Durban JW, Greenfelder M, Guinet C, Jorgensen M, Olson PA, et al. Observations of a distinctive morphotype of killer whale (*Orcinus orca*), type D, from subantarctic waters. *Polar Biol.* 2011;34: 303–306.  
[View Article](#) • [Google Scholar](#)
7. Pitman RL, Durban JW. Killer whale predation on penguins in Antarctica. *Polar Biol.* 2010;33: 1589–1594.  
[View Article](#) • [Google Scholar](#)
8. Pitman RL, Durban JW. Cooperative hunting behavior, prey selectivity and prey handling by pack ice killer whales (*Orcinus orca*), type B, in Antarctica Peninsula waters. *Mar Mamm Sci.* 2012;28: 16–36.  
[View Article](#) • [Google Scholar](#)
9. Morin PA, Archer FI, Foote AD, Vilstrup J, Allen EE, Wade P, et al. Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. *Genome Res.* 2010;20: 908–916. pmid:20413674  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
10. Vongraven D, Bisther A. Prey switching by killer whales in the north-east Atlantic: observational evidence and experimental insights. *Journal of the Marine Biological Association of the UK* 2014;94: 1357–1365.  
[View Article](#) • [Google Scholar](#)
11. Tixier P, Authier M, Gasco N, Guinet C. Influence of artificial food provisioning from fisheries on killer whale reproductive output. *Anim Conserv.* 2015;18: 207–218.  
[View Article](#) • [Google Scholar](#)
12. Ford JKB, Ellis GM. You are what you eat: foraging specializations and their influence on the social organization and behavior of killer whales. In: Yamagiwa J, Karczmarski L, editors. *Primates and Cetaceans: Field Research and Conservation of Complex Mammalian Species.* Primatology Monographs. 2014. pp. 75–98.
13. Foote AD, Vijay N, Ávila-Arcos MC, Baird RW, Durban JW, Fumagalli M, et al. Genome-culture coevolution promotes rapid divergence of killer whale ecotypes. *Nat Commun.* 2016;7: 11693. pmid:27243207  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
14. Hoelzel AR, Hey J, Dahlheim ME, Nicholson C, Burkanov V, Black N. Evolution of population structure in a highly social top predator, the killer whale. *Mol Biol Evol.* 2007;24: 1407–1415. pmid:17400573  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

15. Foote AD, Newton J, Piertney SB, Willerslev E, Gilbert MTP. Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Mol Ecol.* 2009;18: 5207–5217. pmid:20050301  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
16. Foote AD, Vilstrup JT, De Stephanis R, Verborgh P, Nielsen SCA, Deaville R, et al. Genetic differentiation among North Atlantic killer whale populations. *Mol Ecol.* 2011;20: 629–641. pmid:21241391  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
17. Esteban R, Verborgh P, Gauffier P, Giménez J, Martín V, Pérez-Gil M, Tejedor M, Almunia J, Jepson PD, García-Tíscar S, Barrett-Lennars LG, Guinet C, Foote AD, de Stephanis R. Using a multi-disciplinary approach to identify a critically endangered killer whale management unit. *Ecol Indic.* 2016;66: 291–300.  
[View Article](#) • [Google Scholar](#)
18. Sigurjónsson J, Lyrholm T, Leatherwood S, Jónsson E, Víkingsson G. Photoidentification of killer whales, *Orcinus orca*, off Iceland, 1981 through 1986. *Rit Fisk.* 1988;11: 99–114.  
[View Article](#) • [Google Scholar](#)
19. Samarra FIP, Vighi M, Aguilar A, Víkingsson G. Intra-population variation in isotopic niche in herring-eating killer whales. *Mar Ecol Prog Ser.* 2017;564: 199–210.  
[View Article](#) • [Google Scholar](#)
20. Samarra FIP, Foote AD. Seasonal movements of killer whales between Iceland and Scotland. *Aquat Biol.* 2015;24: 75–79.  
[View Article](#) • [Google Scholar](#)
21. Samarra FIP, Tavares SB, Béseau J, Deecke VB, Fennell A, Miller PJO, Pétursson H, Sigurjónsson J, Víkingsson GA. Movements and site fidelity of killer whales (*Orcinus orca*) relative to seasonal and long-term shifts in herring (*Clupea harengus*) distribution. *Mar Biol.* 2017;164: 159.  
[View Article](#) • [Google Scholar](#)
22. Víkingsson GA. Háhyrningur. In: Hersteinsson P, editor. Íslensk spendýr. Reykjavík: Vaka-Helgafell; 2004. pp 166–171.
23. Rasmussen M. The whaling versus whale-watching debate: The resumption of Icelandic whaling. In: Higham J, Bejder L, Williams R, editors. *Whale-watching: Sustainable Tourism and Ecological Management.* Cambridge: Cambridge University Press; 2014. pp. 81–94.
24. Samarra FIP, Tavares S, Miller PJO, Víkingsson GA. Killer whales of Iceland 2006–2015. Report of the Marine and Freshwater Research Institute; 2017 Report No.: HV2017-005 (ISSN 2298-9137).
25. Anonymous. Æðarkollur í morgunmat. *Morgunblaðið.* 2003 Sep 13; p. 6.
26. Anonymous. Úlfar hafsins á æðarfuglsveiðum. *Morgunblaðið.* 2006 Nov 28; p. 18.
27. Skúlason A. Háhyrningar á fuglaveiðum í Berufirði. *Morgunblaðið.* 2010 Dec 1; p. 9.
28. Jonsgård Å, Lyshoel PB. A contribution to the knowledge of the biology of the killer whales *Orcinus orca* (L.). *Nytt Mag Zool.* 1970;18: 41–48.  
[View Article](#) • [Google Scholar](#)
29. Sæmundsson B. Íslensk dýr II: Spendýrin, Reykjavík; 1932.
30. Anonymous. Háhyrningurinn bjargar lúðunni. *Dagblaðið.* 1978 April 27; 4. Árg. 88. TBL. p. 1.
31. Anonymous. Háhyrningar hirtu allar lúður af línunni. *Sjómannaþlaðið Víkingur.* 1979 Sep 1; vol. 41, p. 39.
32. Anonymous. Háhyrningurinn að éta lúðuskipin út á gaddinn. *Morgunblaðið.* 1992 May 13; Blað B.
33. Anonymous. Lítil friður á grálúðumiðum. *Vísir.* 1970 Aug 4.
34. Anonymous. Grálúðan er djúpsjárviskur. *Þjóðviljinn.* 1971 July 25; 36 Árg. 165. TBL. p. 12

35. Donovan G, Gunnlaugsson T. North Atlantic Sightings Surveys 1987: Preliminary report of the aerial survey off Iceland. Rep Int Whal Comm. 1988;39: 437–441.  
[View Article](#) • [Google Scholar](#)
36. Anonymous. Hrafnreiður drepin í Hvalfirði. Fréttir. 1915 Nov 3; vol. 33, p. 137.
37. Anonymous. Hefur skotið 1409 stóröhvelli. Sjómannablaðið Víkingur. 1997 Dec 1; vol. 59, p. 52–55. (excerpt of book Sjávarniður og sunnanrok by Jón Kr. Gunnarsson)
38. Anonymous. “Það er meiri spennna í hrefnuveiðunum”. Morgunblaðið. 1982 June 6; p. 20.
39. Anonymous. Ágangur háhyrninga talin vera orsök. Tíminn. 1967 Feb 21; vol 51(43), p. 16.
40. Bloch D, Lockyer C. Killer whales (*Orcinus orca*) in Faroese waters. Rit Fisk. 1988;11: 55–64.  
[View Article](#) • [Google Scholar](#)
41. Similå T, Holst JC, Christense I. Occurrence and diet of killer whales in northern Norway: seasonal patterns relative to the distribution and abundance of Norwegian spring-spawning herring. Can J Fish Aquat Sci. 1996;53: 769–779.  
[View Article](#) • [Google Scholar](#)
42. Smith WE. Moulting common eiders devoured by killer whales. Br. Birds 2006;99: 264–268.  
[View Article](#) • [Google Scholar](#)
43. Booth CJ, Ellis PM. Common eiders and common guillemots taken by killer whales. Br Birds 2006;99: 533.  
[View Article](#) • [Google Scholar](#)
44. Christensen I. Spekkhoggeren (*Orcinus orca*) I det nordøstlige Atlanterhav. (The killer whales (*Orcinus orca*) in the northeast Atlantic.) Fisk Hav. 1978; 23–31.  
[View Article](#) • [Google Scholar](#)
45. Christensen I. Killer whales in Norwegian coastal waters. Rep Int Whal Comm. 1982;32: 633672.  
[View Article](#) • [Google Scholar](#)
46. Deecke VB, Nykänen M, Foote AD, Janik VM. Vocal behavior and feeding ecology of killer whales *Orcinus orca* around Shetland, UK. Aquat Biol. 2011;13: 79–88.  
[View Article](#) • [Google Scholar](#)
47. Vester H, Hammerschmidt K. First record of killer whales (*Orcinus orca*) feeding on Atlantic salmon (*Salmo salar*) in northern Norway suggest a multi-prey feeding type. Mar Biodivers Rec. 2013;6: e9.  
[View Article](#) • [Google Scholar](#)
48. Weir CR. Killer whales in British waters. Br Wildl. 2002;14: 106–108.  
[View Article](#) • [Google Scholar](#)
49. Luque PL, Davis CG, Reid DG, Wang J, Pierce GJ. Opportunistic sightings of killer whales from Scottish pelagic trawlers fishing for mackerel and herring off North Scotland (UK) between 2000 and 2006. Aquat Living Resour. 2006;19: 403–410.  
[View Article](#) • [Google Scholar](#)
50. Pedro S, Boba C, Dietz R, Sonne C, Rosing-Asvid A, Hansen M, Provas A, McKinney MA. Blubber-depth distribution and bioaccumulation of PCBs and organochlorine pesticides in Arctic-invading killer whales. Sci Total Environ. 2017;601–602: 237–246. pmid:28554115  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
51. Cosentino AM. First record of Norwegian killer whales attacking and feeding on a harbour porpoise. Mar Biodivers Rec. 2015;8: e108.  
[View Article](#) • [Google Scholar](#)
52. Evans PGH. Killer whales (*Orcinus orca*) in British and Irish waters. Rit Fisk. 1988;11: 42–54.  
[View Article](#) • [Google Scholar](#)

53. Galatius A, Kinze CC. *Lagenorhynchus albirostris* (Cetacea: Delphinidae). Mamm Species. 2016;48: 35–47.  
[View Article](#) • [Google Scholar](#)
54. Robinson KP, Bamford CCG, Airey A, Bean TS, Bird C, Haskins GN, et al. Killer whale (*Orcinus orca*) occurrence in the Moray Firth, Northeast Scotland: incidental sightings, behavioural observations, and photo-identification. Aquat Mamm. 2017;43: 26–32.  
[View Article](#) • [Google Scholar](#)
55. Jonsgård A. A note on the attacking behavior of the killer whale (*Orcinus orca*). Nor Hvalfangst-Tid. 1968;57: 84–85.  
[View Article](#) • [Google Scholar](#)
56. Jourdain E, Vongraven D, Bisther A, Karoliussen R. First longitudinal study of seal-feeding killer whales (*Orcinus orca*) in Norwegian coastal waters. PLoS One. 2017;12(6): e0180099. pmid:28666015  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
57. Bolt HE, Harvey PV, Mandleberg L, Foote AD. Occurrence of killer whales in Scottish inshore waters: temporal and spatial patterns relative to the distribution of declining harbor seal populations. Aquat Conserv. 2009;19: 671–675.  
[View Article](#) • [Google Scholar](#)
58. Garðarsson A. Fjöldi æðarfugls, hávellu, toppandar og stökkandar á grunnsævi að vetri. (Numbers of common eider, long-tailed duck, red-breasted merganser, and mallard, wintering on the coast of Iceland). Bliki. 2009;30: 49–54. (in Icelandic)  
[View Article](#) • [Google Scholar](#)
59. Williams AJ, Dyer BM, Randall RM, Komen J. Killer whales *Orcinus orca* and seabirds: “play”, predation and association. Mar Ornithol 1990;18: 37–41.  
[View Article](#) • [Google Scholar](#)
60. Hanson MB, Walker WA. Trans-Pacific consumption of cephalopods by North Pacific killer whales (*Orcinus orca*). Aquat Mamm. 2014;40: 274–284.  
[View Article](#) • [Google Scholar](#)
61. Kennedy J, Jónsson SP, Kasper JM, Ólafsson HG. Movements of female lumpfish (*Cyclopterus lumpus*) around Iceland. ICES J Mar Sci. 2014;72: 880–889.  
[View Article](#) • [Google Scholar](#)
62. Martin AR, Clarke MR. The diet of sperm whales (*Physeter Macrocephalus*) captured between Iceland and Greenland. J Mar Biol Assoc UK 1986;66: 779–790.  
[View Article](#) • [Google Scholar](#)
63. Óskarsson GJ, Taggart CT. Spawning time variation in Icelandic summer-spawning herring (*Clupea harengus*). Can J Fish Aquat Sci. 2009;66: 1666–1681.  
[View Article](#) • [Google Scholar](#)
64. Jónsson B, Jóhannsson M. Fiskirannsóknir á vatnasvæði Þjórsár árið 2013. Report from the Institute of Freshwater Fisheries; 2014 Report No.:VMST/14001.
65. Ford JKB, Ellis GM. Selective foraging by fish-eating killer whales *Orcinus orca* in British Columbia. Mar Ecol Prog Ser. 2006;316: 185–199.  
[View Article](#) • [Google Scholar](#)
66. Anonymous. Tjaldur veiðir gráluðu í net. Morgunblaðið. 2002 Oct 15; vol. 90, issue 241, p. 8.
67. MFRI (2017) Assessment of halibut. Marine and Freshwater Research Institute; 2017 June 13. Available at: [https://www.hafogvatn.is/static/extras/images/Halibut\\_21193.pdf](https://www.hafogvatn.is/static/extras/images/Halibut_21193.pdf)
68. Solmundsson J. Trophic ecology of Greenland halibut (*Reinhardtius hippoglossoides*) on the Icelandic continental shelf and slope. Mar Biol Res. 2007;3: 231–242.  
[View Article](#) • [Google Scholar](#)
69. ICES. Report of the North Western Working Group (NWWG), 27 April—4 May 2017, Copenhagen, Denmark; 2007 Report No.: ICES CM 2017/ACOM:08.

70. De Stephanis R, Giménez J, Esteban R, Gauffier P, García-Tiscar S, Sinding M-HS, et al. Mobbing-like behavior by pilot whales towards killer whales: a response to resource competition or perceived predation risk? *Acta Ethol.* 2016;18: 69–78.  
[View Article](#) • [Google Scholar](#)
71. Stenersen J, Similä T. *Norwegian killer whales*. Henningsvær, Norway: Tringa forlag; 2004.
72. Ford JKB, Ellis GM, Matkin DR, Balcomb KC, Briggs D, Morton AB. Killer whale attacks on minke whales: prey capture and antipredator tactics. *Mar Mamm Sci.* 2005;21: 603–618.  
[View Article](#) • [Google Scholar](#)
73. Bertulli CG, Cecchetti A, Van Bressemer MF, Van Waerebeek K. Skin disorders in common minke whales and white-beaked dolphins off Iceland, a photographic assessment. *J Mar Anim Ecol.* 2012;5: 29–40.  
[View Article](#) • [Google Scholar](#)
74. Gilles A, Gunnlaugsson T, Mikkelsen B, Pike DG, Víkingsson GA. Harbour porpoise *Phocoena phocoena* summer abundance in Icelandic and adjacent waters, based on aerial surveys in 2007 and 2010. *NAMMCO Rep.* 2011;SC/18/AESP/11.  
[View Article](#) • [Google Scholar](#)
75. Bertulli CG, Rasmussen MH, Rosso M. An assessment of the natural marking patterns used for photo-identification of common minke whales and white-beaked dolphins in Icelandic waters. *J Mar Biol Assoc UK* 2016;96: 807–819.  
[View Article](#) • [Google Scholar](#)
76. Baird RW, Dill LM, Stacey PJ. Group-size specific foraging efficiency in transient killer whales (*Orcinus orca*) around southern Vancouver Island. In: Abstracts of the third international orca symposium; 1990 March 9–12; Victoria, BC. Royal British Columbia Museum; 1990. pp. 3.
77. Hauksson E. Abundance of grey seals in Icelandic waters, based on trends of pup-counts from aerial surveys. *NAMMCO Sci Publ.* 2007;6: 85–97.  
[View Article](#) • [Google Scholar](#)
78. Þorbjörnsson JG, Hauksson E, Sigurðsson GM, Granquist SM. Aerial census of the Icelandic harbor seal (*Phoca vitulina*) population in 2016: population estimate, trends and current status/Landselstalning 2016: Stofnstærðarmat, sveiflur og ástand stofns. Report of the Marine and Freshwater Research Institute; 2016 Report No.: HV2017-009 (ISSN 2298-9137).
79. McCordic JA, Todd SK, Stevick PT. Differential rates of killer whale attacks on humpback whales in the North Atlantic as determined by scarification. *J Mar Biol Assoc UK* 2014;94: 1311–1315.  
[View Article](#) • [Google Scholar](#)
80. Mehta AV, Allen JM, Constantine R, Garrigue C, Jann B, Jenner C, et al. Baleen whales are not important as prey for killer whales *Orcinus orca* in high-latitude regions. *Mar Ecol Prog Ser.* 2007;348: 297–307.  
[View Article](#) • [Google Scholar](#)
81. Pitman RL, Totterdell JA, Fearnbach H, Balance LT, Durban JW, Kemps H. Whale killers: prevalence and ecological implications of killer whale predation on humpback whale calves off Western Australia. *Mar Mamm Sci.* 2015;31: 629–657.  
[View Article](#) • [Google Scholar](#)
82. Baird RW, McSweeney DJ, Bane C, Barlow J, Salden DR, Antoine LRK, et al. Killer whales in Hawaiian waters: information on population identity and feeding habits. *Pac Sci.* 2006;60: 523–530.  
[View Article](#) • [Google Scholar](#)
83. Bolaños-Jiménez J, Mignucci-Giannoni AA, Blumenthal J, Bogomolni A, Casas JJ, Henríquez A, et al. Distribution, feeding habits and morphology of killer whales *Orcinus orca* in the Caribbean Sea. *Mamm Rev* 2014;44: 177–189.  
[View Article](#) • [Google Scholar](#)
84. Whitt AD, Baran MA, Bryson M, Rendell LE. First report of killer whales harassing sperm whales in the Gulf of Mexico. *Aquat Mamm.* 2015;41: 252–255.  
[View Article](#) • [Google Scholar](#)



## Publications

[PLOS Biology](#)  
[PLOS Medicine](#)  
[PLOS Computational Biology](#)  
[PLOS Genetics](#)  
[PLOS Pathogens](#)  
[PLOS ONE](#)  
[PLOS Neglected Tropical Diseases](#)

[Home](#)  
[Blogs](#)  
[Collections](#)  
[Give feedback](#)  
[LOCKSS](#)

[Privacy Policy](#)  
[Terms of Use](#)  
[Advertise](#)  
[Media Inquiries](#)  
[Contact](#)

Orcas – *Orcinus orca*. Where do they live? Habitat and distribution. The orca distribution is incredibly broad, ranging from the polar waters to the equator in the open ocean or closed systems such as the Gulf of Mexico, the Gulf of California, the Persian Gulf, the Mediterranean Sea and the Red Sea. They are distributed in all the oceans of the world although they have a preference for cold regions. They have several hunting strategies depending on the type of prey and the region they inhabit. For example, in the North Pacific, they feed on some fish, seabirds, and sharks. *Killer Whales: The Natural History and Genealogy of Orcinus Orca in British Columbia and Washington*. UBC Press, 2000. Killer whales (*Orcinus orca*, Linnaeus, 1758), as a species, have been reported to predate on numerous prey including fish, marine mammals, cephalopods, marine turtles and birds [1, 2]. However, several populations specialize on particular prey types and ecotypes diverging in diet, morphology, genetics and behaviour have been recognised in the North Pacific and Antarctica (e.g. [3–9]). There are numerous observations of killer whales occurring in seasonal herring grounds, making this by far the prey that killer whales are most commonly observed consuming in Iceland. The first published scientific book mentioning herring as a main prey of killer whales dates back to 1932 and since then there have been numerous studies reporting herring consumption (Table 1, #8). Though called killer whales, orcas are not whales. They are the largest dolphin species. In Argentina, orcas hurl themselves onshore to grab sea lion pups. In the Antarctic, pod members cooperate to make large waves that wash seals from ice floes. Whalers call the orca the “killer of whales.” It preys on sperm, gray, fin, humpback, and other whales. 1. 2. 3. 4. WATCH: Looking for Killer Whales 26 Years After the Exxon Valdez Oil Spill. National Geographic travels to Prince William Sound in Alaska to meet with researchers who are studying the lingering effects of the catastrophic Exxon Valdez oil spill. Killer whales have a cosmopolitan distribution and as a species are generalists, feeding on a variety of prey. However, local populations tend to specialise on specific prey types. In Icelandic waters, killer whales are generally associated with herring and, thus, have been presumed to be herring specialists. However, recent studies suggest a more complex foraging ecology, possibly including a mosaic of strategies. With increased observational effort in recent years due to research and whale-watching activities, there have been several reports of interactions with different prey, including con Fast Facts: Killer Whales (Orcas). Scientific Name: *Orcinus orca*. Common Name(s): Killer whale, orca, blackfish, grampus. Basic Animal Group: Mammal. Size: 16–26 feet. Killer whales may work in pods to hunt their prey and have a number of interesting techniques to hunt prey, which includes working together to create waves to wash seals off ice floes and sliding onto beaches to capture prey. Killer whales use a variety of sounds for communicating, socializing and finding prey. Determining species of killer whales is important not only in gaining information about the whales but in protecting them—it is difficult to determine the abundance of killer whales without even knowing how many species there are. Killer Whales and Humans.